

PAINLESS LEFT HEMORRHAGIC PLEURAL EFFUSION: AN UNUSUAL PRESENTATION OF LEAKING SACCULAR AORTIC ARCH ANEURYSM

Mohammed Azfar Siddiqui,¹ Jamal Akhtar,² Syed Wajahat Ali Rizvi,³
Syed Amjad Ali Rizvi,⁴ Kaleem Ahmad,⁵ Ibne Ahmad,¹ Ekramullah¹

¹ Department of Radiodiagnosis, Jawaharlal Nehru Medical College, A.M.U. Aligarh, India

² Department of Tuberculosis and Chest Disease, Jawaharlal Nehru Medical College, A.M.U. Aligarh, India

³ Department of Ophthalmology, Jawaharlal Nehru Medical College, A.M.U. Aligarh, India

⁴ Department of Surgery, Jawaharlal Nehru Medical College, A.M.U. Aligarh, India

⁵ Department of Radiodiagnosis, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

PJR July - September 2011; 21(3): 123-126

ABSTRACT

Most thoracic aortic aneurysms are asymptomatic and are detected by chance on routine chest imaging for some other reasons. Only rarely it is symptomatic due to leak and dissection which is a potentially life threatening event that commonly presents with severe pain. In this report, we present the case of a 67-year-old man who presented with shortness of breath, intermittent cough, fever, and left sided painless hemorrhagic pleural effusion. Further investigation by plain radiography, CT and MRI revealed a saccular aneurysm arising from the lateral aspect of the mid-transverse arch of the aorta, along with a dissecting descending aortic aneurysm with false lumen communicating with left pleural space. The patient refused any surgical procedure and was treated conservatively with blood transfusions and anti hypertensive medication. On the 8th day patient finally succumb to a fatal episode of shock. We suggest dissecting thoracic aneurysm be included in the differential diagnosis of nontraumatic hemorrhagic pleural effusion in an elderly patient presenting with dysnea, cough and fever, which otherwise suggest the clinical diagnosis of bronchogenic carcinoma. CT of the chest should be immediately performed as the diagnostic procedure of choice.

Key words: Aneurysm; Aorta; Saccular; Dissecting; Leaking; Hemorrhagic; Pleural effusion

Case Report

A 67-year-old known hypertensive man was admitted to the Chest Medicine Department with complaints of new onset shortness of breath, intermittent cough, and fever without chills. There was no history of sputum, hemoptysis, and chest pain or weight loss. The patient also denied any history of trauma, cyanosis, jaundice or any secondary complication of hypertension. His blood pressure was 156/92 mm Hg, and pulse rate was 68 beats/min. On physical examination there was dullness to percussion and reduced breath sounds at left lung base. There was also decreased movement

and diminished vocal fremitus on left side. Cardiac examination did not reveal any abnormalities and his electrocardiogram was also unremarkable.

Biochemical analysis demonstrated normal cardiac enzyme level and presence of normochromic and normocytic anemia with hemoglobin of 10 g/100 mL. Chest X-ray was done, which shows massive left sided effusion and an abnormal convex opacity in the aortopulmonary area with right mediastinal shift (Fig. 1-A.). The patient was put in a propped up bed with moist O₂ and IV fluid with Cephalosporin and Deriphylline was started.

A left thoracocentesis was performed, and it revealed thin, grossly hemorrhagic, reddish pleural fluid that did

Correspondence : Dr. Mohammed Azfar Siddiqui
Department of Radiodiagnosis
Jawaharlal Nehru Medical College, AMU
Aligarh-202002, India
Tel: 0091-9719361454 (M)
Email: drazfarsiddiqui@gmail.com

not clot. Fluid cytology and biochemistry shows plenty of RBCs with elevated neutrophil count. No organisms were identified on Gram's stain or culture, nor were malignant cells identified by cytology.




Figure 1-A: Frontal scout CT image shows massive left sided pleural effusion and an abnormal convex opacity in the aortopulmonary area with right mediastinal shift. **Figure 1-B:** CT axial image at the level of aortic arch shows the presence of a saccular aneurysm arising from the lateral aspect of the mid-transverse arch of the aorta, along with a left sided pleural effusion.

Computed Tomography (CT) scan revealed a saccular aneurysm arising from the lateral aspect of the mid-transverse arch of the aorta, along with a dissecting descending aortic aneurysm (Fig. 1 & 2). An intimal flap was seen separating the true lumina and false

lumina, which communicated with the left pleural space. The presence of hyperdense fluid in the left pleural space signified a leaking aortic aneurysm. Magnetic Resonance Imaging (MRI) was done which confirm the nature of lesion (Fig. 3).



Figure 2-A: Axial CT image at the level of left atrium shows dissecting descending aortic aneurysm with false lumen communicating with pleural space along with leaking hyperdense hemorrhagic fluid in pleural space. **Figure 2-B:** Sagittal reconstructed CT image shows the presence of saccular aneurysm and dissecting descending aortic aneurysm.



Figure 3-A: Coronal T2W TrueFISP MR image demonstrated a saccular aneurysm arising from aortic arch with variable signal intensity fluid in left pleural space signifying hemorrhage in various stages. **Figure 3-B:** Time of flight MR Angiography again demonstrated a saccular aneurysm arising from aortic arch.

Two units of packed RBCs were administered and the patient was treated with propranolol. Cardiothoracic surgery consultation was done which emphasized need for aggressive preoperative management of arterial pressure and prompt surgical intervention. However, patient refused any surgical procedures. On the 8th day of admission, patient condition deteriorated and all resuscitative measures failed with patient dying within 4 hours of beginning of fatal episode of shock.

Discussion

Aneurysm is a localized or diffuse dilation of an artery with a diameter at least 50% greater than the expected size of the artery.¹ In the ascending aorta, a diameter larger than 4 cm while in the descending aorta, a diameter greater than 3 cm is regarded as an aneurysm.² In a true aneurysm, all of the components of vessel wall are present, whereas a false aneurysm has an incomplete wall. The shape of aneurysm may be fusiform (involve the entire circumference of the aortic wall) or saccular (involve only a portion of the wall).

Aortic dissection is characterized by dissection of blood along the laminar planes of the aortic media, with the formation of a blood-filled channel within the aortic wall. It commonly occurs in two groups of patients.³ The first group consists of older men with long history of hypertension. The second major subgroup consists of younger patients with a systemic or localized abnormality of aortic connective tissue that include aortic coarctation, a bicuspid aortic valve, and disorders of collagen, including Marfan's syndrome, Ehlers-Danlos syndrome, and degeneration of the aortic media.

Most thoracic aortic aneurysms are asymptomatic and are detected by chance on chest X-ray.⁴ Few cases present with sudden death due to acute rupture. In patients who survive the initial tear, severe pain is a classic presenting symptom. Pain is tearing in nature and located either in the anterior chest, which is suggestive of an ascending aortic dissection, or in the posterior chest or back, which is suggestive of a descending aortic dissection.³ Compressive symptoms like hoarseness of voice, stridor, cough, wheeze, left diaphragmatic palsy, dysphagia, are also reported as presenting symptoms.⁴ Painless dissection was described only in a minority of the patients, and most of these presented with congestive heart failure, stroke, or syncope.⁵

Hemorrhagic pleural effusion may occur in various conditions. The different diagnosis includes traumatic injury, pulmonary infarction, tuberculosis, pulmonary thromboembolism, and pleuropulmonary malignancy. It could also be a presenting sign of acute aortic dissection that is extremely rare.⁶ Besides our case, only few cases have been reported in literature with such presentation.

Acute aortic dissection is a potentially life-threatening condition requiring immediate assessment. Prompt

and accurate diagnosis is required to initiate appropriate surgical repair or medical treatment. Without immediate treatment, the outcome is often fatal with more than 50% of patients dying in the first 48 hours.⁶ However, misdiagnosis still remains an unresolved problem because of variable and unpredictable clinical presentation.

In the past, angiography was the only accurate examination for evaluating the aorta. Currently, noninvasive radiologic assessment of patients with techniques such as spiral CT, MRI, and Transesophageal Echocardiography (TEE) is the cornerstone of the diagnostic process.⁷ A chest x-ray is usually the initial examination performed and reveals pathologic findings such as abnormal aortic contour, widening of mediastinum, displaced intimal calcification, and pleural effusion.⁸ A contrast-enhanced CT scan currently is the method of choice for the diagnosis and management of patients with suspected dissection because of a diagnostic accuracy comparable with aortography, wide availability, ease of performance, and examination speed.⁷ Due to continued improvements in technique, MRI and TEE is also being performed in increasing numbers of patients with suspected aortic dissection. Angiographic evaluation is now reserved for only those patients in whom the previously noted studies are equivocal or when additional anatomic information is required especially for those in whom surgery is planned.⁷ Based on current recommendations most of the uncomplicated descending aortic dissection are treated medically. Surgical approach is reserved for patients with proximal dissection or for cases of distal dissection complicated by rupture, compromise of a major vessel, or recurrent pain.^{7,9,10}

In summary, our case highlights the unpredictable clinical presentation of a potentially life threatening condition, dissecting aortic aneurysm. Although in an elderly male who present with X-ray evidences of massive pleural effusion and hilar mass, bronchogenic carcinoma comes as a most important differential diagnosis, but this case study suggest that aortic dissection, though rare should be kept in mind as a differential diagnosis in these situations.

References

1. Benjamin ME, Hansen KJ, Craven TE, Keith DR, Plonk GW, Geary RL, Dean RH. Combined aortic and renal artery surgery. A contemporary experience. *Ann Surg.* 1996 May; **223**(5): 555-65.
2. Erbel R. Diseases of the thoracic aorta. *Heart.* 2001 Aug; **86**(2): 227-34.
3. Little S, Johnson J, Moon BY, Mehta S. Painless left hemorrhagic pleural effusion: An unusual presentation of dissecting ascending aortic aneurysm. *Chest.* 1999 Nov; **116**(5): 1478-80.
4. Sengupta P, Mitra B, Saha K, Maitra S, Pal J, Sarkar N. Descending thoracic aortic aneurysm presenting as left sided hemorrhagic pleural effusion. *J Assoc Physicians India.* 2007 Apr; **55**: 297-300.
5. Spittell PC, Spittell JA Jr, Joyce JW, Tajik AJ, Edwards WD, Schaff HV, Stanson AW. Clinical features and differential diagnosis of aortic dissection: experience with 236 cases (1980 through 1990). *Mayo Clin Proc.* 1993 Jul; **68**(7):642-51.
6. Chen HY. Rupture of aortic aneurysm with late right hemorrhagic pleural effusion. *Am J Emerg Med.* 2010 Nov; **28**(9): 1058.e1-3.
7. Gandelman G, Barzilay N, Krupsky M, Resnitzky P. Left pleural hemorrhagic effusion. A presenting sign of thoracic aortic dissecting aneurysm. *Chest.* 1994 Aug; **106**(2):636-8.
8. Slater EE, DeSanctis RW. The clinical recognition of dissecting aortic aneurysm. *Am J Med.* 1976 May 10; **60**(5): 625-33.
9. Glower DD, Speier RH, White WD, Smith LR, Rankin JS, Wolfe WG. Management and long-term outcome of aortic dissection. *Ann Surg.* 1991 Jul; **214**(1): 31-41.
10. Glower DD, Fann JI, Speier RH, Morrison L, White WD, Smith LR, Rankin JS, Miller DC, Wolfe WG. Comparison of medical and surgical therapy for uncomplicated descending aortic dissection. *Circulation.* 1990 Nov; **82**(5): 39-46.